

# 19 Transit time crossing a river

**Lab Objective:** *This lab discusses a classical calculus of variations problem: how is a river to be crossed in the shortest possible time? We will look at a numerical solution using the pseudospectral method.*

Suppose a boat is to be rowed across a river, from a point  $A$  on one side of a river ( $x = -1$ ), to a point  $B$  on the other side ( $x = 1$ ). Assuming the boat moves at a constant speed 1 relative to the current, how must the boat be steered to minimize the time required to cross the river?

Let us consider a typical trajectory for the boat as it crosses the river. If  $T$  is the time required to cross the river, then the position  $s$  of the boat at time  $t$  is

$$\begin{aligned} s(t) &= \langle x(t), y(t) \rangle, \quad t \in [0, T], \\ s'(t) &= \langle x'(t), y'(t) \rangle, \\ &= \langle \cos \theta(x(t)), \sin \theta(x(t)) \rangle + \langle 0, c(x(t)) \rangle. \end{aligned}$$

Here  $\langle \cos \theta, \sin \theta \rangle$  represents the motion of the boat due to the rower, and  $\langle 0, c \rangle$  is the motion of the boat due to the current.

We can relate the angle at which the boat is steered to the graph of its trajectory by noting that

$$\begin{aligned} y'(x) &= \frac{y'(t)}{x'(t)}, \\ &= \frac{\sin \theta + c}{\cos \theta}, \\ &= c \sec \theta + \tan \theta. \end{aligned} \tag{19.1}$$

The time  $T$  required to cross the river is given by

$$\begin{aligned} T &= \int_{-1}^1 t'(x) dx, \\ &= \int_{-1}^1 \frac{1}{x'(t)} dx \\ &= \int_{-1}^1 \sec(\theta) dx. \end{aligned} \tag{19.2}$$




Figure 19.1: The river's current, along with a possible trajectory for the boat.

We would like to find an expression for the total time  $T$  required to cross the river from  $A$  to  $B$ , in terms of the graph of the boat's trajectory. To derive the functional  $T[y]$ , we note that

$$\begin{aligned} T[y] &= \int_{-1}^1 \sec \theta \, dx, \\ &= \int_{-1}^1 \frac{1}{1-c^2} (c \tan \theta + \sec \theta - c^2 \sec \theta - c \tan \theta) \, dx, \\ &= \int_{-1}^1 \frac{1}{1-c^2} (c \tan \theta + \sec \theta - cy') \, dx. \end{aligned}$$

Since

$$\begin{aligned} c \tan \theta + \sec \theta &= \sqrt{1-c^2 + (c \sec \theta + \tan \theta)^2}, \\ &= \sqrt{1-c^2 + (y')^2}, \end{aligned}$$

we obtain at last

$$T[y] = \int_{-1}^1 \left[ \alpha(x) \sqrt{1 + (\alpha y')^2(x)} - (\alpha^2 c y')(x) \right] dx, \quad (19.3)$$

where  $\alpha = (1-c^2)^{-1/2}$ .

**Problem 1.** Assume that the current is given by  $c(x) = -\frac{7}{10}(x^2 - 1)$ . (This function assumes, for example, that the current is faster near the center of the river.) Write 2 python functions. The first should accept as arguments a function  $y$ , its derivative  $y'$ , and an  $x$ -value, and return  $L(x, y(x), y'(x))$  (where  $T[y] = \int_{-1}^1 L(x, y(x), y'(x))$ ). The second should use the first function to compute and return  $T[y]$  for a given path  $y(x)$ . (Hint: The integration for  $T[y]$  can be done use an approximation method such as the midpoint method or can be done using the `quad` function from `scipy.integrate`.)

**Problem 2.** Let  $y(x)$  be the straight-line path between  $A = (-1, 0)$  and  $B = (1, 5)$ . Numerically calculate  $T[y]$  to get an upper bound on the minimum time required to cross from  $A$  to  $B$ . Using (19.2), find a lower bound on the minimum time required to cross. (Hint: if  $G = \int f(x)dx$  and we want to minimize  $G$ , try minimizing  $f(x)$ .)

We look for the path  $y(x)$  that minimizes the time required for the boat to cross the river, so that the function  $T$  is minimized. From the calculus of variations we know that a smooth path  $y(x)$  minimizes  $T$  only if the Euler-Lagrange equation is satisfied. Recall that the Euler-Lagrange equation is

$$L_y - \frac{d}{dx}L_{y'} = 0.$$

Since  $L_y = 0$ , we see that the shortest time trajectory satisfies

$$\frac{d}{dx}L_{y'} = \frac{d}{dx} \left( \alpha^3(x)y'(x)(1 + (\alpha y')^2(x))^{-1/2} - \alpha^2(x)c \right) = 0. \quad (19.4)$$

**Problem 3.** Numerically solve the Euler-Lagrange equation (19.4), using  $c(x) = -\frac{7}{10}(x^2 - 1)$  and  $\alpha = (1 - c^2)^{-1/2}$ , and  $y(-1) = 0$ ,  $y(1) = 5$ .

Hint: Since this boundary value problem is defined over the domain  $[-1, 1]$ , it is easy to solve using the pseudospectral method. Begin by replacing each  $\frac{d}{dx}$  with the pseudospectral differentiation matrix  $D$ . Then impose the boundary conditions and solve.

**Problem 4.** Plot the angle at which the boat should be pointed at each  $x$ -coordinate. (Hint: Use Equation (19.1); see Figure 19.3. Note that the angle the boat should be steered is *not* described by the tangent vector to the trajectory. Consider using `scipy.optimize.root` or `scipy.interpolate.barycentric_interpolate`)




Figure 19.2: Numerical computation of the trajectory with the shortest transit time.




Figure 19.3: The optimal angle to steer the boat.